
High-resolution, data consistent modelling of the Alps' last glaciation coupled with 3D particle tracking achieved with physics-informed machine learning

Tancrede Leger^{*1,2}, Guillaume Jovet , Sarah Kamleitner , Jürgen Mey , Frédéric Herman , Brandon Finley , Maxime Bernard , Balthazar Allegri , Susan Ivy-Ochs , Andreas Vieli , Andreas Henz , and Samuel Nussbaumer

¹University of Sheffield [Sheffield] – Western Bank Sheffield S10 2TN, Royaume-Uni

²Université de Lausanne = University of Lausanne (UNIL) – CH-1015 Lausanne, Suisse

Résumé

Reconstructing the last glaciation of the European Alpine Ice Field via numerical modelling has been challenged by persistent model-data disagreements, including large overestimations of its former thickness. Here, we tackle this issue by applying the Instructed Glacier Model, a three-dimensional, high-order, and thermo-mechanically coupled model enhanced with physics-informed machine learning. This new approach allows us to produce an ensemble of 100, Alps-wide and 22 thousand-year-long (40-18 ka) simulations at 300 m spatial resolution. Unfeasible with traditional models due to computational costs, our experiment substantially increases model-data agreement in both ice extent and thickness. The offset in ice thickness, for instance, is here reduced by between 200% and 450% relative to previous studies. In a second experiment, we leverage the parallelization of GPU computing to couple, for the first time, our Alpine Ice Field model with 3D and time-transgressive ice advection of particles (tens of millions). Here, particles are seeded to mimic both the subglacial (e.g. abrasion, plucking) and supraglacial (e.g. rockfall) origins of glacial sediments. Using our ensemble best-fit simulation, we present the results of tracking the sink-to-source transport trajectories of distinct LGM ice-contact deposits (e.g. terminal moraines), and the LGM source-to-sink transport trajectories of specific surface lithologies, across the Alps. We find that modelling the Alps-wide transport of glacial sediments also helps us better understand the complex internal ice dynamics of the former Alpine Ice Field, including former ice transfluences and the zipping/unzipping behaviours of different tributary glaciers. More generally, this work demonstrates that physics-informed AI-driven glacier models can overcome the bottleneck of high-resolution continental-scale modelling required to accurately describe complex topographies and glacier dynamics.

Mots-Clés: Quaternary glaciations, Alpine Ice Field, Last Glacial maximum, Numerical modelling, machine learning, particle tracking

*Intervenant